



WE CLAIM:


1. A process for making a hydroxamic acid compound or a salt thereof, wherein:
the process comprises reacting a 4-sulfonyloxy-heterocyclyl compound or a 4-
5 halo-heterocyclyl compound with a metal thioester; and
the hydroxamic acid compound corresponds in structure to Formula (1-1):




- the 4-sulfonyloxy-heterocyclyl compound corresponds in structure to Formula (1-
2):



- the 4-halo-heterocyclyl compound corresponds in structure to Formula (1-3):



- the metal thioester corresponds in structure to Formula (1-4):



- n is selected from the group consisting of zero, 1, and 2; and

X¹ is selected from the group consisting of -O-, -S-, -S(O)-, -S(O)₂-, and -N(R^{x1})-;
and

X⁴ is selected from the group consisting of alkyl, haloalkyl, aryl, and haloaryl; and

X⁵ is selected from the group consisting of alkyl, aryl, and arylalkyl, wherein:

5 the alkyl, aryl, and arylalkyl are optionally substituted with one or more
independently selected halogen; and

X⁷ is halogen; and

M is a metal cation; and

Z is selected from the group consisting of -O-, -S-, -S(O)-, -S(O)₂-, and -N(R^x)-;

10 and

Z³ is selected from the group consisting of nitrogen and carbon bonded to
hydrogen; and

R^x is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, R^a-
oxyalkyl, aminosulfonyl, alkylsulfonyl, R^aR^a-aminoalkyl, carbocyclyl, carbocyclylalkyl,
15 carbocyclylsulfonyl, heterocyclyl, heterocyclylalkyl, and heterocyclylsulfonyl, wherein:

any member of such group optionally is substituted with one or more
substituents independently selected from the group consisting of halogen, hydroxy,
cyano, amino, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, imino, alkyl,
alkoxy, alkylthio, alkoxyalkyl, and alkoxyalkoxy, wherein:

20 any member of such group optionally is substituted with one or
more substituents independently selected from the group consisting of
halogen, hydroxy, and alkyl; and

R^{x1} is a nitrogen-protecting group; and

A is selected from the group consisting of -O-, -S-, -S(O)-, -S(O)₂-, -C(O)-, -NR^b-,
25 -CO-N(R^b), -N(R^b)-C(O)-, -C(O)-O-, -O-C(O)-, -O-C(O)-O-, -HC=CH-, -C≡C-, -N=N-,
-C(S)-N(R^b)-, -N(R^b)-C(S)-, alkyl, alkoxy, oxyalkyl, alkylthio, thioalkyl, and a bond; and

R is selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxyalkyl,
carbocyclyl, heterocyclyl, carbocyclylalkyl, heterocyclylalkyl, carbocyclylalkoxyalkyl,
heterocyclylalkoxyalkyl, carbocyclylthioalkyl, and heterocyclylthioalkyl, wherein:

30 any member of such group optionally is substituted with one or more
substituents independently selected from the group consisting of halogen, nitro,

nitroso, hydroxy, oxo, alkyl, alkoxy, alkylthio, alkoxyalkyl, alkoxyalkylalkyl, C₁-C₂-alkylenedioxy, and alkoxyalkyl, wherein:

5 any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino; and

E is selected from the group consisting of -O-, -C(O)-, -C(O)-O-, -O-C(O)-, -N(R^b)-, -C(O)-N(R^b)-, -N(R^b)-C(O)-, -C(O)-N(R^b)-N(R^b)-C(O)-, -N(R^b)-C(O)-N(R^b)-, -S-, -S(O)-, -S(O)₂-, -N(R^b)-S(O)₂-, -S(O)₂-N(R^b)-, -O-S(O)₂-, -S(O)₂-O-, -C(NH)-, 10 -C(NOH)-, -N(R^b)-C(NH)-, -N(R^b)-C(NOH)-, -C(NH)-N(R^b)-, -C(NOH)-N(R^b)-, alkyl, alkenyl, carbonylalkyl, alkylcarbonyl, and a bond, wherein:

any alkyl or alkenyl portion of a substituent in such group optionally is substituted with one or more independently selected R^c substituents; and

Y is selected from the group consisting of hydrogen, halogen, cyano, alkyl, 15 alkenyl, alkynyl, alkoxyalkyl, alkoxyalkoxyalkyl, alkylthioalkyl, alkylthioalkylthioalkyl, alkylthioalkoxyalkyl, alkoxyalkylthioalkyl, carbocyclyl, carbocyclylalkyl, carbocyclylalkoxyalkyl, heterocyclyl, heterocyclylalkyl, and heterocyclylalkoxyalkyl, wherein:

20 any member of such group optionally is substituted with one or more independently selected R^d substituents; and

each R^a is independently selected from the group consisting of hydrogen, hydroxy, alkyl, alkenyl, alkynyl, alkoxy, alkoxyalkyl, bisalkoxyalkyl, alkylthioalkyl, alkylthioalkenyl, alkylsulfoxidoalkyl, alkylsulfonyl, alkylsulfonylalkyl, carbocyclyl, carbocyclylalkyl, carbocyclylalkoxyalkyl, carbocyclylthioalkyl, 25 carbocyclylthioalkenyl, carbocyclylsulfoxidoalkyl, carbocyclylsulfonyl, carbocyclylsulfonylalkyl, heterocyclyl, heterocyclylalkyl, heterocyclylalkoxyalkyl, heterocyclylthioalkyl, heterocyclylsulfoxidoalkyl, heterocyclylsulfonyl, heterocyclylsulfonylalkyl, aminoalkyl, aminosulfonyl, aminoalkylsulfonyl, and alkoxyalkylaminoalkyl, wherein:

30 any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, imino, alkyl, alkylcarbonyl, carbocyclyl, and carbocyclylalkyl; and

each R^b is independently selected aryl; and

each R^c is independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, -C(H)(NH), -C(H)(NOH), thiol, sulfo, nitro, nitroso, oxo, thioxo, imino, alkyl, alkoxy, alkenyl, alkynyl, alkoxyalkyl, alkylthio, carbocyclyl, carbocyclylalkyl,

5 heterocyclyl, and heterocyclylalkyl, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, imino, alkyl, and carbocyclylalkyl; and

10 each R^d is independently selected from the group consisting of halogen, hydroxy, cyano, sulfo, nitro, nitroso, oxo, thioxo, imino, alkyl, alkoxy, alkoxyalkyl, -C(O)(R^e), -S-R^e, -S(O)₂-R^e, carbocyclyl, alkylcarbocyclyl, carbocyclylalkyl, heterocyclyl, alkylheterocyclyl, and heterocyclylalkyl, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino; and

15 each R^e is independently selected from the group consisting of hydrogen alkyl, carbocyclyl, carbocyclylalkyl, heterocyclyl, and heterocyclylalkyl, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino; and

20 each R^f is independently selected from the group consisting of hydrogen, alkyl, -O-R^h, carbocyclylalkyl, and heterocyclylalkyl; and



any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino; and

25 each R^h is independently selected from the group consisting of hydrogen, alkyl, carbocyclyl, carbocyclylalkyl, heterocyclyl, and heterocyclylalkyl, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino.

2. A process according to claim 1, wherein:

the hydroxamic acid compound corresponds in structure to a formula selected from the group consisting of:



5 R^2 is selected from the group consisting of aryl and heteroaryl, wherein:


the aryl or heteroaryl optionally is substituted with one or more substituents independently selected from the group consisting of halogen, nitro, nitroso, hydroxy, alkyl, alkoxy, alkylthio, alkoxyalkyl, alkoxyalkylalkyl, C_1 - C_2 -alkylenedioxy, and alkoxyalkyl, wherein:

10 any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino.

15 3. A process according to claim 2, wherein Z is $-N(R^x)-$:


4. A process according to claim 2, wherein the metal thioester comprises potassium thioacetate.

20 5. A process according to claim 2, wherein the 4-sulfonyloxy-heterocyclyl compound corresponds in structure to Formula (5-1):




(5-1)

and



(5-2).

6. A process according to claim 2, wherein the hydroxamic acid compound corresponds in structure to Formula (6-1):



(6-1).


5

7. A process according to claim 2, wherein the process comprises reacting the 4-halo-heterocyclyl compound with the metal thioester.


8. A process according to claim 2, wherein the process comprises reacting the
10 4-sulfonyloxy-heterocyclyl compound with the metal thioester.

9. A process according to claim 8, wherein:
the process comprises:

15 reacting the 4-sulfonyloxy-heterocyclyl compound with the metal
thioester to form a 4-thioester-heterocyclyl compound, and
oxidatively halogenating the 4-thioester-heterocyclyl compound to
form a 4-halosulfonyl-heterocyclyl compound; and
the 4-thioester-heterocyclyl compound corresponds in structure to Formula (9-1):



the 4-halosulfonyl-heterocyclyl compound corresponds in structure to Formula (9-2):



5 X^2 is halogen.

10. A process according to claim 9, wherein:

X^2 is chloro; and

the oxidative halogenation comprises combining the 4-thioester-heterocyclyl
10 compound with a source of Cl_2 , N-chlorosuccinimide, or 1,3-dichloro-5,5-dimethylhydantoin.

11. A process according to claim 9, wherein X^1 is $-N(R^{x1})-$.


15 12. A process according to claim 11, wherein:

the process further comprises:


reacting a second amount of the 4-sulfonyloxy-heterocyclyl
compound with an alcohol to form a protected cyclic amino compound
having a nitrogen protecting group (R^{x1}), and

20 removing the nitrogen protecting group from the cyclic amino
compound to form an unprotected cyclic amino compound, and

reacting the 4-halosulfonyl-heterocyclcyl compound with the
unprotected cyclic amino compound; and
the alcohol is HO-R-E-Y; and
the protected cyclic amino compound corresponds in structure to Formula (12-1):




the unprotected cyclic amino compound corresponds in structure to Formula (12-2):




- 10 13. A process according to claim 12, wherein:
R is phenyl, and
E-Y is selected from the group consisting of fluoroalkyl and fluoroalkoxy.


14. A process according to claim 13, wherein the hydroxamic acid compound
15 corresponds in structure to Formula (14-1):




and




15. A process for making a 4-thioester-heterocyclcyl compound, wherein:
the process comprises reacting a 4-sulfonyloxy-heterocyclcyl compound or a 4-
20 halo-heterocyclcyl compound with a metal thioester; and
the 4-thioester-heterocyclcyl compound corresponds in structure to Formula (15-1):




the 4-sulfonyloxy-heterocyclyl compound corresponds in structure to Formula (15-2):



5 the 4-halo-heterocyclyl compound corresponds in structure to Formula (15-3):



the metal thioester corresponds in structure to Formula (15-4):



X^1 is selected from the group consisting of -O-, -S-, -S(O)-, -S(O)₂-, and -N(R^{x1})-;

10 and

X^4 is selected from the group consisting of alkyl, haloalkyl, aryl, and haloaryl; and

X^5 is selected from the group consisting of alkyl, aryl, and arylalkyl, wherein:


the alkyl, aryl, and arylalkyl are optionally substituted with one or more independently selected halogen; and

X^7 is halogen; and


M is a metal cation; and

R^{x1} is a nitrogen-protecting group.

- 5 16. A process according to claim 15, wherein the 4-thioester-heterocyclyl compound corresponds in structure to Formula (16-1):




and



17. A compound or a salt thereof, wherein:

the compound corresponds in structure to Formula (17-1):



10

X^1 is selected from the group consisting of -O-, -S-, -S(O)-, -S(O)₂-, and -N(R^{x1})-;

and

X^5 is selected from the group consisting of alkyl, aryl, and arylalkyl, wherein:


the alkyl, aryl, and arylalkyl are optionally substituted with one or more

15

independently selected halogen; and


R^{x1} is a nitrogen-protecting group.

18. A compound or salt thereof according to claim 17, wherein the compound corresponds in structure to Formula (18-1):



(18-1)


and



(18-2).


19. A process for making a hydroxamic acid compound or a salt thereof, wherein:
the process comprises oxidatively halogenating a 4-thioester-heterocyclyl
compound; and

5 the hydroxamic acid compound corresponds in structure to Formula (19-1):



(19-1); and

the 4-thioester-heterocyclyl compound corresponds in structure to Formula (19-2):



(19-2); and

n is selected from the group consisting of zero, 1, and 2; and

10 X¹ is selected from the group consisting of -O-, -S-, -S(O)-, -S(O)₂-, and -N(R^{x1})-;

and

X⁵ is selected from the group consisting of alkyl, aryl, and arylalkyl, wherein:

the alkyl, aryl, and arylalkyl are optionally substituted with one or more
independently selected halogen; and

15 Z is selected from the group consisting of -O-, -S-, -S(O)-, -S(O)₂-, and -N(R^x)-;

and

Z^3 is selected from the group consisting of nitrogen and carbon bonded to hydrogen; and

R^x is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, R^a -oxyalkyl, aminosulfonyl, alkylsulfonyl, R^aR^a -aminoalkyl, carbocyclyl, carbocyclylalkyl, carbocyclylsulfonyl, heterocyclyl, heterocyclylalkyl, and heterocyclylsulfonyl, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, amino, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, imino, alkyl, alkoxy, alkylthio, alkoxyalkyl, and alkoxyalkoxy, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, and alkyl; and

R^{x1} is a nitrogen-protecting group; and

A is selected from the group consisting of -O-, -S-, -S(O)-, -S(O)₂-, -C(O)-, -NR^b-, -CO-N(R^b), -N(R^b)-C(O)-, -C(O)-O-, -O-C(O)-, -O-C(O)-O-, -HC=CH-, -C≡C-, -N=N-, -C(S)-N(R^b)-, -N(R^b)-C(S)-, alkyl, alkoxy, oxyalkyl, alkylthio, thioalkyl, and a bond; and

R is selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxyalkyl, carbocyclyl, heterocyclyl, carbocyclylalkyl, heterocyclylalkyl, carbocycliloxyalkyl, heterocycliloxyalkyl, carbocyclylthioalkyl, and heterocyclylthioalkyl, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, nitro, nitroso, hydroxy, oxo, alkyl, alkoxy, alkylthio, alkoxyalkyl, alkoxyalkylalkyl, C₁-C₂-alkylenedioxy, and alkoxyalkyl, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino; and

E is selected from the group consisting of -O-, -C(O)-, -C(O)-O-, -O-C(O)-, -N(R^b)-, -C(O)-N(R^b)-, -N(R^b)-C(O)-, -C(O)-N(R^b)-N(R^b)-C(O)-, -N(R^b)-C(O)-N(R^b)-, -S-, -S(O)-, -S(O)₂-, -N(R^b)-S(O)₂-, -S(O)₂-N(R^b)-, -O-S(O)₂-, -S(O)₂-O-, -C(NH)-, -C(NOH)-, -N(R^b)-C(NH)-, -N(R^b)-C(NOH)-, -C(NH)-N(R^b)-, -C(NOH)-N(R^b)-, alkyl, alkenyl, carbonylalkyl, alkylcarbonyl, and a bond, wherein:

any alkyl or alkenyl portion of a substituent in such group optionally is substituted with one or more independently selected R^c substituents; and

Y is selected from the group consisting of hydrogen, halogen, cyano, alkyl, alkenyl, alkynyl, alkoxyalkyl, alkoxyalkoxyalkyl, alkylthioalkyl, alkylthioalkylthioalkyl, alkylthioalkoxyalkyl, alkoxyalkylthioalkyl, carbocyclyl, carbocyclylalkyl, carbocyclylalkoxyalkyl, heterocyclyl, heterocyclylalkyl, and heterocyclylalkoxyalkyl, wherein:

any member of such group optionally is substituted with one or more independently selected R^d substituents; and

each R^a is independently selected from the group consisting of hydrogen, hydroxy, alkyl, alkenyl, alkynyl, alkoxy, alkoxyalkyl, bisalkoxyalkyl, alkylthioalkyl, alkylthioalkenyl, alkylsulfoxidoalkyl, alkylsulfonyl, alkylsulfonylalkyl, carbocyclyl, carbocyclylalkyl, carbocyclylalkoxyalkyl, carbocyclylthioalkyl, carbocyclylthioalkenyl, carbocyclylsulfoxidoalkyl, carbocyclylsulfonyl, carbocyclylsulfonylalkyl, heterocyclyl, heterocyclylalkyl, heterocyclylalkoxyalkyl, heterocyclylthioalkyl, heterocyclylsulfoxidoalkyl, heterocyclylsulfonyl, heterocyclylsulfonylalkyl, aminoalkyl, aminosulfonyl, aminoalkylsulfonyl, and alkoxyalkylaminoalkyl, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, imino, alkyl, alkylcarbonyl, carbocyclyl, and carbocyclylalkyl; and

each R^b is independently selected aryl; and

each R^c is independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, -C(H)(NH), -C(H)(NOH), thiol, sulfo, nitro, nitroso, oxo, thioxo, imino, alkyl, alkoxy, alkenyl, alkynyl, alkoxyalkyl, alkylthio, carbocyclyl, carbocyclylalkyl, heterocyclyl, and heterocyclylalkyl, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, imino, alkyl, and carbocyclylalkyl; and

each R^d is independently selected from the group consisting of halogen, hydroxy, cyano, sulfo, nitro, nitroso, oxo, thioxo, imino, alkyl, alkoxy, alkoxyalkyl, -C(O)(R^g), -S-

R^e , $-S(O)_2-R^e$, carbocyclyl, alkylcarbocyclyl, carbocyclylalkyl, heterocyclyl, alkylheterocyclyl, and heterocyclylalkyl, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino; and each R^e is independently selected from the group consisting of hydrogen alkyl, carbocyclyl, carbocyclylalkyl, heterocyclyl, and heterocyclylalkyl, wherein:


any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino; and each R^g is independently selected from the group consisting of hydrogen, alkyl, $-O-R^h$, carbocyclylalkyl, and heterocyclylalkyl; and


any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino; and each R^h is independently selected from the group consisting of hydrogen, alkyl, carbocyclyl, carbocyclylalkyl, heterocyclyl, and heterocyclylalkyl, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino.

20. A process according to claim 19, wherein R^{x1} is selected from the group consisting of alkoxyalkyl, alkoxycarbonyl, and arylalkoxycarbonyl.

21. A process according to claim 19, wherein:
the hydroxamic acid compound corresponds in structure to a formula selected from the group consisting of:





R^2 is selected from the group consisting of aryl and heteroaryl, wherein:

the aryl or heteroaryl optionally is substituted with one or more substituents independently selected from the group consisting of halogen, nitro, nitroso, hydroxy, alkyl, alkoxy, alkylthio, alkoxyalkyl, alkoxycarbonylalkyl, C_1 - C_2 -alkylenedioxy, and alkoxycarbonyl, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino.

22. A process according to claim 21, wherein Z is $-N(R^x)-$.

23. A process according to claim 21, wherein the oxidative halogenation comprises combining the 4-thioester-heterocyclyl compound with a source of Cl_2 , N-chlorosuccinimide, or 1,3-dichloro-5,5-dimethylhydantoin.

24. A process according to claim 23, wherein the combination of the 4-thioester-heterocyclyl compound with the source of Cl_2 , N-chlorosuccinimide, or 1,3-dichloro-5,5-dimethylhydantoin occurs in the presence of an alcohol, acetic acid, or a chloroalkyl solvent.

25. A process according to claim 24, wherein the 4-thioester-heterocyclyl compound is combined with a source of Cl_2 .


26. A process according to claim 25, wherein the combination of the 4-thioester-heterocyclyl compound with the source of Cl_2 occurs in the presence of ethanol.

27. A process according to claim 25, wherein the combination of the 4-thioester-heterocyclyl compound with the source of Cl_2 occurs in the presence of glacial acetic acid.

28. A process according to claim 25, wherein the combination of the 4-thioester-heterocyclyl compound with the source of Cl_2 occurs in the presence of a chloroalkyl solvent selected from the group consisting of CCl_4 , CHCl_3 , and CCl_2 .

29. A process according to claim 24, wherein:

the 4-thioester-heterocyclyl compound corresponds in structure to Formula (29-1):




(29-1), and

the 4-thioester-heterocyclyl compound is combined with a source of Cl_2 in the presence of ethanol.

30. A process according to claim 24, wherein:


the 4-thioester-heterocyclyl compound corresponds in structure to Formula (30-1):



(30-1), and


the 4-thioester-heterocyclyl compound is combined with a source of Cl_2 in the presence of glacial acetic acid.

31. A process according to claim 24, wherein the hydroxamic acid compound corresponds in structure to Formula (31-1):




(31-1).

32. A process for making a 4-halosulfonyl-heterocyclyl compound, wherein:
the process comprises oxidatively halogenating a 4-thioester-heterocyclyl
5 compound; and
the 4-halosulfonyl-heterocyclyl compound corresponds in structure to Formula
(32-1):



(32-1); and

the 4-thioester-heterocyclyl compound corresponds in structure to Formula (32-2):



(32-2); and


X¹ is selected from the group consisting of -O-, -S-, -S(O)-, -S(O)₂-, and -N(R^{x1})-;
and

X² is halogen; and

X⁵ is selected from the group consisting of alkyl, aryl, and arylalkyl, wherein:

the alkyl, aryl, and arylalkyl are optionally substituted with one or more independently selected halogen; and
 R^{x1} is a nitrogen-protecting group.


- 5 33. A compound or a salt thereof, wherein:
the compound corresponds in structure to Formula (33-1):




X^1 is selected from the group consisting of -O-, -S-, -S(O)-, -S(O)₂-, and -N(R^{x1})-;
and

- 10 X^2 is halogen; and
 R^{x1} is a nitrogen-protecting group.


34. A compound or salt thereof according to claim 33, wherein the compound corresponds in structure to Formula (34-1):



and




- 15 35. A process for making a hydroxamic acid compound or a salt thereof, wherein:
the process comprises reacting a 4-halosulfur-heterocyclyl compound with a cyclic amino compound; and
the hydroxamic acid compound corresponds in structure to Formula (35-1):




(35-1); and

the 4-halosulfur-heterocyclyl compound corresponds in structure to Formula (35-2):



(35-2); and

the cyclic amino compound corresponds in structure to Formula (35-3):



(35-3); and

each n is independently selected from the group consisting of zero, 1, and 2; and X¹ is selected from the group consisting of -O-, -S-, -S(O)-, -S(O)₂-, and -N(R^{x1})-

and

X² is halogen; and

Z is selected from the group consisting of -O-, -S-, -S(O)-, -S(O)₂-, and -N(R^x)-

and

Z³ is selected from the group consisting of nitrogen and carbon bonded to hydrogen; and

R^x is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, R^a-oxyalkyl, aminosulfonyl, alkylsulfonyl, R^aR^a-aminoalkyl, carbocyclyl, carbocyclylalkyl, carbocyclylsulfonyl, heterocyclyl, heterocyclylalkyl, and heterocyclylsulfonyl, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, amino, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, imino, alkyl, alkoxy, alkylthio, alkoxyalkyl, and alkoxyalkoxy, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, and alkyl; and

R^{x1} is a nitrogen-protecting group; and

5 A is selected from the group consisting of -O-, -S-, -S(O)-, -S(O)₂-, -C(O)-, -NR^b-, -CO-N(R^b), -N(R^b)-C(O)-, -C(O)-O-, -O-C(O)-, -O-C(O)-O-, -HC=CH-, -C≡C-, -N=N-, -C(S)-N(R^b)-, -N(R^b)-C(S)-, alkyl, alkoxy, oxyalkyl, alkylthio, thioalkyl, and a bond; and

R is selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxyalkyl, carbocyclyl, heterocyclyl, carbocyclylalkyl, heterocyclylalkyl, carbocycliloxyalkyl, heterocycliloxyalkyl, carbocyclylthioalkyl, and heterocyclylthioalkyl, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, nitro, nitroso, hydroxy, oxo, alkyl, alkoxy, alkylthio, alkoxyalkyl, alkoxycarbonylalkyl, C₁-C₂-alkylenedioxy, and alkoxycarbonyl, wherein:

15 any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino; and

E is selected from the group consisting of -O-, -C(O)-, -C(O)-O-, -O-C(O)-, -N(R^b)-, -C(O)-N(R^b)-, -N(R^b)-C(O)-, -C(O)-N(R^b)-N(R^b)-C(O)-, -N(R^b)-C(O)-N(R^b)-, -S-, -S(O)-, -S(O)₂-, -N(R^b)-S(O)₂-, -S(O)₂-N(R^b)-, -O-S(O)₂-, -S(O)₂-O-, -C(NH)-, -C(NOH)-, -N(R^b)-C(NH)-, -N(R^b)-C(NOH)-, -C(NH)-N(R^b)-, -C(NOH)-N(R^b)-, alkyl, alkenyl, carbonylalkyl, alkylcarbonyl, and a bond, wherein:

any alkyl or alkenyl portion of a substituent in such group optionally is substituted with one or more independently selected R^c substituents; and

25 Y is selected from the group consisting of hydrogen, halogen, cyano, alkyl, alkenyl, alkynyl, alkoxyalkyl, alkoxyalkoxyalkyl, alkylthioalkyl, alkylthioalkylthioalkyl, alkylthioalkoxyalkyl, alkoxyalkylthioalkyl, carbocyclyl, carbocyclylalkyl, carbocyclylalkoxyalkyl, heterocyclyl, heterocyclylalkyl, and heterocyclylalkoxyalkyl, wherein:

any member of such group optionally is substituted with one or more independently selected R^d substituents; and

each R^a is independently selected from the group consisting of hydrogen, hydroxy, alkyl, alkenyl, alkynyl, alkoxy, alkoxyalkyl, bisalkoxyalkyl, alkylthioalkyl, alkylthioalkenyl, alkylsulfoxidoalkyl, alkylsulfonyl, alkylsulfonylalkyl, carbocyclyl, carbocyclylalkyl, carbocyclyloxyalkyl, carbocyclylalkoxyalkyl, carbocyclylthioalkyl, carbocyclylthioalkenyl, carbocyclylsulfoxidoalkyl, carbocyclylsulfonyl, carbocyclylsulfonylalkyl, heterocyclyl, heterocyclylalkyl, heterocyclyloxyalkyl, heterocyclylalkoxyalkyl, heterocyclylthioalkyl, heterocyclylsulfoxidoalkyl, heterocyclylsulfonyl, heterocyclylsulfonylalkyl, aminoalkyl, aminosulfonyl, aminoalkylsulfonyl, and alkoxyalkylaminoalkyl, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, imino, alkyl, alkylcarbonyl, carbocyclyl, and carbocyclylalkyl; and

each R^b is independently selected aryl; and

each R^c is independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, -C(H)(NH), -C(H)(NOH), thiol, sulfo, nitro, nitroso, oxo, thioxo, imino, alkyl, alkoxy, alkenyl, alkynyl, alkoxyalkyl, alkylthio, carbocyclyl, carbocyclylalkyl, heterocyclyl, and heterocyclylalkyl, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, imino, alkyl, and carbocyclylalkyl; and

each R^d is independently selected from the group consisting of halogen, hydroxy, cyano, sulfo, nitro, nitroso, oxo, thioxo, imino, alkyl, alkoxy, alkoxyalkyl, -C(O)(R^e), -S-R^e, -S(O)₂-R^e, carbocyclyl, alkylcarbocyclyl, carbocyclylalkyl, heterocyclyl, alkylheterocyclyl, and heterocyclylalkyl, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino; and

each R^e is independently selected from the group consisting of hydrogen alkyl, carbocyclyl, carbocyclylalkyl, heterocyclyl, and heterocyclylalkyl, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino; and

each R^g is independently selected from the group consisting of hydrogen, alkyl, -

5 O- R^h , carbocyclalkyl, and heterocyclalkyl; and

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino; and

each R^h is independently selected from the group consisting of hydrogen, alkyl,

10 carbocycl, carbocyclalkyl, heterocycl, and heterocyclalkyl, wherein:



any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino.

15 36. A process according to claim 35, wherein R^{x1} is selected from the group consisting of alkoxyalkyl, alkoxycarbonyl, and arylalkoxycarbonyl.



37. A process according to claim 35, wherein:

the hydroxamic acid compound corresponds in structure to a formula selected from

20 the group consisting of:



the cyclic amino compound corresponds in structure to a formula selected from the group consisting of:



R² is selected from the group consisting of aryl and heteroaryl, wherein:

the aryl or heteroaryl optionally is substituted with one or more substituents independently selected from the group consisting of halogen, nitro, nitroso, hydroxy, alkyl, alkoxy, alkylthio, alkoxyalkyl, alkoxycarbonylalkyl, C₁-C₂-alkylenedioxy, and alkoxycarbonyl, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino.

38. A process according to claim 37, wherein:

R and R² are optionally-substituted aryl; and

A is selected from the group consisting of -O-, -S-, -S(O)₂-, -O-S(O)₂-, -S(O)₂-O-, -C(O)-, -C(O)-O-, -O-C(O)-, and a bond; and




E is selected from the group consisting of -O-, -C(O)-, and a bond.

39. A process according to claim 38, wherein:


X¹ is -N(R^{x1})-; and

R^{x1} is selected from the group consisting of alkoxyalkyl, alkoxycarbonyl, and arylalkoxycarbonyl.

40. A process according to claim 39, wherein the 4-halosulfur-heterocyclyl compound corresponds in structure to Formula (40-1):



41. A process according to claim 37, wherein the hydroxamic acid compound corresponds in structure to Formula (41-1):



5

42. A process according to claim 38, wherein:

the cyclic amino compound is prepared by a process comprising a reaction wherein a reagent is 4-hydroxy-piperidine, and


the 4-halosulfur-heterocyclyl compound is prepared by a process comprising a
10 reaction wherein a reagent is 4-hydroxy-piperidine.

43. A process according to claim 38, wherein:

the cyclic amino compound is prepared by a process comprising a reaction wherein a reagent is a first amount of a protected 4-hydroxy-piperidinyl compound; and

5 the 4-halosulfur-heterocyclyl compound is prepared by a process comprising a reaction wherein a reagent is a second amount of the protected 4-hydroxy-piperidinyl compound; and

the protected 4-hydroxy-piperidinyl compound corresponds in structure to Formula (43-1):



10

(43-1); and


R^{x1} is selected from the group consisting of alkoxycarbonyl and arylalkoxycarbonyl.

44. A process according to claim 38, wherein:

15 the cyclic amino compound is prepared by a process comprising a reaction wherein a reagent is a first amount of a 4-sulfonyloxy-piperidinyl compound; and

the 4-halosulfur-heterocyclyl compound is prepared by a process comprising a reaction wherein a reagent is a second amount of the 4-sulfonyloxy-piperidinyl compound; and

20 the 4-sulfonyloxy-piperidinyl compound corresponds in structure to Formula (44-1):



(44-1); and

R^{x1} is selected from the group consisting of alkoxycarbonyl and arylalkoxycarbonyl; and

X^4 is selected from the group consisting of alkyl, haloalkyl, aryl, and haloaryl.

45. A process according to claim 44, wherein:


the cyclic amino compound is prepared by a process comprising:

reacting the 4-sulfonyloxy-heterocyclyl compound with an alcohol to form a protected cyclic amino compound having a nitrogen protecting group (R^{x1}), and

removing the nitrogen protecting group from the protected cyclic amino compound; and

the alcohol is $HO-R-E-Y$; and

the protected cyclic amino compound corresponds in structure to Formula (45-1):




46. A process according to claim 44, wherein:

the 4-halosulfur-heterocyclyl compound is prepared by a process comprising:


reacting the 4-sulfonyloxy-heterocyclyl compound with a metal thioester to form a 4-thioester-heterocyclyl compound, and

oxidatively halogenating the 4-thioester-heterocyclyl compound; and

the metal thioester corresponds in structure to Formula (46-1):



the 4-thioester-heterocyclyl compound corresponds in structure to Formula (46-2):




M is a metal cation; and


X⁵ is selected from the group consisting of alkyl, aryl, and arylalkyl, wherein:
the alkyl, aryl, and arylalkyl are optionally substituted with one or more
independently selected halogen.

47. A process according to claim 46, wherein the oxidative halogenation comprises combining the 4-thioester-heterocyclyl compound with a source of Cl₂, N-chlorosuccinimide, or 1,3-dichloro-5,5-dimethylhydantoin.


48. A process for making an sulfuramine compound, wherein:
the process comprises reacting a 4-halosulfur-heterocyclyl compound with a cyclic amino compound; and
the sulfuramine compound corresponds in structure to Formula (48-1):



the 4-halosulfur-heterocyclyl compound corresponds in structure to Formula (48-2):



the cyclic amino compound corresponds in structure to Formula (48-3):



each n is independently selected from the group consisting of zero, 1, and 2; and

Z³ is selected from the group consisting of nitrogen and carbon bonded to

5 hydrogen; and

X¹ is selected from the group consisting of -O-, -S-, -S(O)-, -S(O)₂-, and -N(R^{x1})-;

and

X² is halogen; and

R^{x1} is a nitrogen-protecting group; and

10 A is selected from the group consisting of -O-, -S-, -S(O)-, -S(O)₂-, -C(O)-, -NR^b-,
-CO-N(R^b), -N(R^b)-C(O)-, -C(O)-O-, -O-C(O)-, -O-C(O)-O-, -HC=CH-, -C≡C-, -N=N-,
-C(S)-N(R^b)-, -N(R^b)-C(S)-, alkyl, alkoxy, oxyalkyl, alkylthio, thioalkyl, and a bond; and

R is selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxyalkyl,
carbocyclyl, heterocyclyl, carbocyclylalkyl, heterocyclylalkyl, carbocycloxyalkyl,

15 heterocycloxyalkyl, carbocyclylthioalkyl, and heterocyclylthioalkyl, wherein:

any member of such group optionally is substituted with one or more
substituents independently selected from the group consisting of halogen, nitro,
nitroso, hydroxy, oxo, alkyl, alkoxy, alkylthio, alkoxyalkyl, alkoxycarbonylalkyl,
C₁-C₂-alkylenedioxy, and alkoxycarbonyl, wherein:

20 any member of such group optionally is substituted with one or
more substituents independently selected from the group consisting of
halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo,
and imino; and

E is selected from the group consisting of -O-, -C(O)-, -C(O)-O-, -O-C(O)-,
25 -N(R^b)-, -C(O)-N(R^b)-, -N(R^b)-C(O)-, -C(O)-N(R^b)-N(R^b)-C(O)-, -N(R^b)-C(O)-N(R^b)-,
-S-, -S(O)-, -S(O)₂-, -N(R^b)-S(O)₂-, -S(O)₂-N(R^b)-, -O-S(O)₂-, -S(O)₂-O-, -C(NH)-,
-C(NOH)-, -N(R^b)-C(NH)-, -N(R^b)-C(NOH)-, -C(NH)-N(R^b)-, -C(NOH)-N(R^b)-, alkyl,
alkenyl, carbonylalkyl, alkylcarbonyl, and a bond, wherein:

any alkyl or alkenyl portion of a substituent in such group optionally is
30 substituted with one or more independently selected R^c substituents; and

Y is selected from the group consisting of hydrogen, halogen, cyano, alkyl, alkenyl, alkynyl, alkoxyalkyl, alkoxyalkoxyalkyl, alkylthioalkyl, alkylthioalkylthioalkyl, alkylthioalkoxyalkyl, alkoxyalkylthioalkyl, carbocyclyl, carbocyclylalkyl, carbocyclylalkoxyalkyl, heterocyclyl, heterocyclylalkyl, and heterocyclylalkoxyalkyl,
5 wherein:

any member of such group optionally is substituted with one or more independently selected R^d substituents; and
each R^b is independently selected aryl; and

each R^c is independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, -C(H)(NH), -C(H)(NOH), thiol, sulfo, nitro, nitroso, oxo, thioxo, imino, alkyl, alkoxy, alkenyl, alkynyl, alkoxyalkyl, alkylthio, carbocyclyl, carbocyclylalkyl, heterocyclyl, and heterocyclylalkyl, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, imino, alkyl, and carbocyclylalkyl; and

each R^d is independently selected from the group consisting of halogen, hydroxy, cyano, sulfo, nitro, nitroso, oxo, thioxo, imino, alkyl, alkoxy, alkoxyalkyl, -C(O)(R^g), -S-R^e, -S(O)₂-R^e, carbocyclyl, alkylcarbocyclyl, carbocyclylalkyl, heterocyclyl, alkylheterocyclyl, and heterocyclylalkyl, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino; and
each R^e is independently selected from the group consisting of hydrogen alkyl, carbocyclyl, carbocyclylalkyl, heterocyclyl, and heterocyclylalkyl, wherein:


any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino; and
each R^g is independently selected from the group consisting of hydrogen, alkyl, -O-R^h, carbocyclylalkyl, and heterocyclylalkyl; and

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino; and

each R^h is independently selected from the group consisting of hydrogen, alkyl, carbocyclyl, carbocyclylalkyl, heterocyclyl, and heterocyclylalkyl, wherein:


any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino.

49. A process according to claim 48, wherein the sulfuramine compound corresponds in structure to Formula (49-1):



(49-1)


and



(49-2).


50. A process for making a hydroxamic acid compound or a salt thereof, wherein:

the process comprises removing a nitrogen-protecting group (R^{x1}) from a piperidinyll nitrogen of a nitrogen-protected piperidinyll compound; and the hydroxamic acid compound corresponds in structure to Formula (50-1):



(50-1); and

the nitrogen-protected piperidinyll compound corresponds in structure to Formula (50-2):



Z^3 is selected from the group consisting of nitrogen and carbon bonded to hydrogen; and

each n is independently selected from the group consisting of zero, 1, and 2; and

5 R^x is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, R^a -oxyalkyl, alkylsulfonyl, aminosulfonyl, R^aR^a -aminoalkyl, carbocyclyl, carbocyclylalkyl, carbocyclylsulfonyl, heterocyclyl, heterocyclylalkyl, and heterocyclylsulfonyl, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, amino, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, imino, alkyl, 10 alkoxy, alkylthio, alkoxyalkyl, and alkoxyalkoxy, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, and alkyl; and

15 A is selected from the group consisting of $-O-$, $-S-$, $-S(O)-$, $-S(O)_2-$, $-C(O)-$, $-NR^b-$, $-CO-N(R^b)-$, $-N(R^b)-C(O)-$, $-C(O)-O-$, $-O-C(O)-$, $-O-C(O)-O-$, $-HC=CH-$, $-C\equiv C-$, $-N=N-$, $-C(S)-N(R^b)-$, $-N(R^b)-C(S)-$, alkyl, alkoxy, oxyalkyl, alkylthio, thioalkyl, and a bond; and

R is selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxyalkyl, carbocyclyl, heterocyclyl, carbocyclylalkyl, heterocyclylalkyl, carbocycliloxyalkyl, 20 heterocycliloxyalkyl, carbocyclylthioalkyl, and heterocyclylthioalkyl, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, nitro, nitroso, hydroxy, oxo, alkyl, alkoxy, alkylthio, alkoxyalkyl, alkoxyalkonylalkyl, C_1-C_2 -alkylenedioxy, and alkoxyalkonyl, wherein:

25 any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino; and

E is selected from the group consisting of -O-, -C(O)-, -C(O)-O-, -O-C(O)-, -N(R^b)-, -C(O)-N(R^b)-, -N(R^b)-C(O)-, -C(O)-N(R^b)-N(R^b)-C(O)-, -N(R^b)-C(O)-N(R^b)-, -S-, -S(O)-, -S(O)₂-, -N(R^b)-S(O)₂-, -S(O)₂-N(R^b)-, -O-S(O)₂-, -S(O)₂-O-, -C(NH)-, -C(NOH)-, -N(R^b)-C(NH)-, -N(R^b)-C(NOH)-, -C(NH)-N(R^b)-, -C(NOH)-N(R^b)-, alkyl, alkenyl, carbonylalkyl, alkylcarbonyl, and a bond, wherein:

any alkyl or alkenyl portion of a substituent in such group optionally is substituted with one or more independently selected R^c substituents; and

Y is selected from the group consisting of hydrogen, halogen, cyano, alkyl, alkenyl, alkynyl, alkoxyalkyl, alkoxyalkoxyalkyl, alkylthioalkyl, alkylthioalkylthioalkyl, alkylthioalkoxyalkyl, alkoxyalkylthioalkyl, carbocyclyl, carbocyclylalkyl, carbocyclylalkoxyalkyl, heterocyclyl, heterocyclylalkyl, and heterocyclylalkoxyalkyl, wherein:

any member of such group optionally is substituted with one or more independently selected R^d substituents; and

each R^a is independently selected from the group consisting of hydrogen, hydroxy, alkyl, alkenyl, alkynyl, alkoxy, alkoxyalkyl, bisalkoxyalkyl, alkylthioalkyl, alkylthioalkenyl, alkylsulfoxidoalkyl, alkylsulfonyl, alkylsulfonylalkyl, carbocyclyl, carbocyclylalkyl, carbocyclylalkoxyalkyl, carbocyclylthioalkyl, carbocyclylthioalkenyl, carbocyclylsulfoxidoalkyl, carbocyclylsulfonyl, carbocyclylsulfonylalkyl, heterocyclyl, heterocyclylalkyl, heterocyclylalkoxyalkyl, heterocyclylthioalkyl, heterocyclylsulfoxidoalkyl, heterocyclylsulfonyl, heterocyclylsulfonylalkyl, aminoalkyl, aminosulfonyl, aminoalkylsulfonyl, and alkoxyalkylaminoalkyl, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, imino, alkyl, alkylcarbonyl, carbocyclyl, and carbocyclylalkyl; and

each R^b is independently selected aryl; and

each R^c is independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, -C(H)(NH), -C(H)(NOH), thiol, sulfo, nitro, nitroso, oxo, thioxo, imino, alkyl, alkoxy, alkenyl, alkynyl, alkoxyalkyl, alkylthio, carbocyclyl, carbocyclylalkyl, heterocyclyl, and heterocyclylalkyl, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, imino, alkyl, and carbocyclalkyl; and

5 each R^d is independently selected from the group consisting of halogen, hydroxy, cyano, sulfo, nitro, nitroso, oxo, thioxo, imino, alkyl, alkoxy, alkoxyalkyl, -C(O)(R^e), -S-R^e, -S(O)₂-R^e, carbocycl, alkylcarbocycl, carbocyclalkyl, heterocycl, alkylheterocycl, and heterocyclalkyl, wherein:

10 any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino; and each R^e is independently selected from the group consisting of hydrogen alkyl, carbocycl, carbocyclalkyl, heterocycl, and heterocyclalkyl, wherein:

15 any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino; and each R^f is independently selected from the group consisting of hydrogen, alkyl, -O-R^h, carbocyclalkyl, and heterocyclalkyl; and



20 any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino; and each R^h is independently selected from the group consisting of hydrogen, alkyl, carbocycl, carbocyclalkyl, heterocycl, and heterocyclalkyl, wherein:

25 any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino.



51. A process according to claim 50, wherein R^{x1} is selected from the group consisting of alkoxycarbonyl and arylalkoxycarbonyl.

30

52. A process according to claim 50, wherein:
the hydroxamic acid compound corresponds in structure to a formula selected from the group consisting of:



the nitrogen-protected piperidinyll compound corresponds in structure to a formula selected from the group consisting of:



R^2 is selected from the group consisting of aryl and heteroaryl, wherein:

the aryl or heteroaryl optionally is substituted with one or more substituents independently selected from the group consisting of halogen, nitro, nitroso, hydroxy, alkyl, alkoxy, alkylthio, alkoxyalkyl, alkoxyalkyl, C_1 - C_2 -alkylenedioxy, and alkoxyalkyl, wherein:

10

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thio, and imino.

15

53. A process according to claim 52, wherein the process comprises contacting the nitrogen-protected piperidinyl compound with a hydrogen source and a transition metal catalyst.

5 54. A process according to claim 53, wherein the hydrogen source comprises ammonium formate.

55. A process according to claim 52, wherein the process comprises contacting the nitrogen-protected piperidinyl compound with an acid.

10

56. A process according to claim 55, wherein the acid has a pKa of no greater than about -3.

57. A process according to claim 55, wherein contacting of the nitrogen-protected piperidinyl compound with the acid occurs at a temperature of greater than 30°C.


15

58. A process according to claim 56, wherein the acid comprises HCl gas.

59. A process according to claim 56, wherein the acid comprises HCl in an alcohol.


20

60. A process according to claim 55, wherein the nitrogen-protected piperidinyl compound corresponds in structure selected from the group consisting of:




(60-1)

and




(60-2).

25 61. A process according to claim 52, wherein the hydroxamic acid compound corresponds in structure to Formula (61-1):




(61-1).

62. A process for making an unprotected piperidinyl compound, wherein:
the process comprises removing a nitrogen-protecting group (R^{x1}) from a
5 piperidinyl nitrogen of a nitrogen-protected piperidinyl compound; and
the unprotected piperidinyl compound corresponds in structure to Formula (62-1):



(62-1); and

the nitrogen-protected piperidinyl compound corresponds in structure to Formula
(62-2):



(62-2); and

Z^3 is selected from the group consisting of nitrogen and carbon bonded to
hydrogen; and

each n is independently selected from the group consisting of zero, 1, and 2; and

- A is selected from the group consisting of $-\text{O}-$, $-\text{S}-$, $-\text{S}(\text{O})-$, $-\text{S}(\text{O})_2-$, $-\text{C}(\text{O})-$, $-\text{NR}^{\text{b}}-$,
15 $-\text{CO}-\text{N}(\text{R}^{\text{b}})-$, $-\text{N}(\text{R}^{\text{b}})-\text{C}(\text{O})-$, $-\text{C}(\text{O})-\text{O}-$, $-\text{O}-\text{C}(\text{O})-$, $-\text{O}-\text{C}(\text{O})-\text{O}-$, $-\text{HC}=\text{CH}-$, $-\text{C}\equiv\text{C}-$, $-\text{N}=\text{N}-$,
 $-\text{C}(\text{S})-\text{N}(\text{R}^{\text{b}})-$, $-\text{N}(\text{R}^{\text{b}})-\text{C}(\text{S})-$, alkyl, alkoxy, oxyalkyl, alkylthio, thioalkyl, and a bond; and

R is selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxyalkyl,
carbocyclyl, heterocyclyl, carbocyclylalkyl, heterocyclylalkyl, carbocycloxyalkyl,
heterocycloxyalkyl, carbocyclylthioalkyl, and heterocyclylthioalkyl, wherein:

- 20 any member of such group optionally is substituted with one or more
substituents independently selected from the group consisting of halogen, nitro,

nitroso, hydroxy, oxo, alkyl, alkoxy, alkylthio, alkoxyalkyl, alkoxycarbonylalkyl, C₁-C₂-alkylenedioxy, and alkoxycarbonyl, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino; and

E is selected from the group consisting of -O-, -C(O)-, -C(O)-O-, -O-C(O)-, -N(R^b)-, -C(O)-N(R^b)-, -N(R^b)-C(O)-, -C(O)-N(R^b)-N(R^b)-C(O)-, -N(R^b)-C(O)-N(R^b)-, -S-, -S(O)-, -S(O)₂-, -N(R^b)-S(O)₂-, -S(O)₂-N(R^b)-, -O-S(O)₂-, -S(O)₂-O-, -C(NH)-, -C(NOH)-, -N(R^b)-C(NH)-, -N(R^b)-C(NOH)-, -C(NH)-N(R^b)-, -C(NOH)-N(R^b)-, alkyl, alkenyl, carbonylalkyl, alkylcarbonyl, and a bond, wherein:

any alkyl or alkenyl portion of a substituent in such group optionally is substituted with one or more independently selected R^c substituents; and

Y is selected from the group consisting of hydrogen, halogen, cyano, alkyl, alkenyl, alkynyl, alkoxyalkyl, alkoxyalkoxyalkyl, alkylthioalkyl, alkylthioalkylthioalkyl, alkylthioalkoxyalkyl, alkoxyalkylthioalkyl, carbocyclyl, carbocyclylalkyl, carbocyclylalkoxyalkyl, heterocyclyl, heterocyclylalkyl, and heterocyclylalkoxyalkyl, wherein:

any member of such group optionally is substituted with one or more independently selected R^d substituents; and

each R^b is independently selected aryl; and

each R^c is independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, -C(H)(NH), -C(H)(NOH), thiol, sulfo, nitro, nitroso, oxo, thioxo, imino, alkyl, alkoxy, alkenyl, alkynyl, alkoxyalkyl, alkylthio, carbocyclyl, carbocyclylalkyl, heterocyclyl, and heterocyclylalkyl, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, imino, alkyl, and carbocyclylalkyl; and

each R^d is independently selected from the group consisting of halogen, hydroxy, cyano, sulfo, nitro, nitroso, oxo, thioxo, imino, alkyl, alkoxy, alkoxyalkyl, -C(O)(R^g), -S-R^e, -S(O)₂-R^e, carbocyclyl, alkylcarbocyclyl, carbocyclylalkyl, heterocyclyl, alkylheterocyclyl, and heterocyclylalkyl, wherein:


any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino; and each R^e is independently selected from the group consisting of hydrogen alkyl, carbocyclyl, carbocyclylalkyl, heterocyclyl, and heterocyclylalkyl, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino; and each R^g is independently selected from the group consisting of hydrogen, alkyl, -O-R^h, carbocyclylalkyl, and heterocyclylalkyl; and

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino; and each R^h is independently selected from the group consisting of hydrogen, alkyl, carbocyclyl, carbocyclylalkyl, heterocyclyl, and heterocyclylalkyl, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino.


63. A process according to claim 62, wherein the unprotected piperidinyl compound corresponds in structure to Formula (63-1):



64. A process for making a hydroxamic acid compound or a salt thereof, wherein:


the process comprises contacting an unprotected piperidinyl compound with an N-alkylating agent; and

the hydroxamic acid compound corresponds in structure to Formula (64-1):



(64-1); and

the unprotected piperidiny compound corresponds in structure to Formula (64-2):



(64-2); and

Z^3 is selected from the group consisting of nitrogen and carbon bonded to

5 hydrogen; and

each n is independently selected from the group consisting of zero, 1, and 2; and

A is selected from the group consisting of $-O-$, $-S-$, $-S(O)-$, $-S(O)_2-$, $-C(O)-$, $-NR^b-$, $-CO-N(R^b)-$, $-N(R^b)-C(O)-$, $-C(O)-O-$, $-O-C(O)-$, $-O-C(O)-O-$, $-HC=CH-$, $-C\equiv C-$, $-N=N-$, $-C(S)-N(R^b)-$, $-N(R^b)-C(S)-$, alkyl, alkoxy, oxyalkyl, alkylthio, thioalkyl, and a bond; and

10 R is selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxyalkyl, carbocyclyl, heterocyclyl, carbocyclylalkyl, heterocyclylalkyl, carbocycloxyalkyl, heterocycloxyalkyl, carbocyclylthioalkyl, and heterocyclylthioalkyl, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, nitro, nitroso, hydroxy, oxo, alkyl, alkoxy, alkylthio, alkoxyalkyl, alkoxycarbonylalkyl, C_1-C_2 -alkylenedioxy, and alkoxycarbonyl, wherein:

15

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino; and

20

E is selected from the group consisting of $-O-$, $-C(O)-$, $-C(O)-O-$, $-O-C(O)-$, $-N(R^b)-$, $-C(O)-N(R^b)-$, $-N(R^b)-C(O)-$, $-C(O)-N(R^b)-N(R^b)-C(O)-$, $-N(R^b)-C(O)-N(R^b)-$, $-S-$, $-S(O)-$, $-S(O)_2-$, $-N(R^b)-S(O)_2-$, $-S(O)_2-N(R^b)-$, $-O-S(O)_2-$, $-S(O)_2-O-$, $-C(NH)-$,

-C(NOH)-, -N(R^b)-C(NH)-, -N(R^b)-C(NOH)-, -C(NH)-N(R^b)-, -C(NOH)-N(R^b)-, alkyl, alkenyl, carbonylalkyl, alkylcarbonyl, and a bond, wherein:

any alkyl or alkenyl portion of a substituent in such group optionally is substituted with one or more independently selected R^c substituents; and

5 Y is selected from the group consisting of hydrogen, halogen, cyano, alkyl, alkenyl, alkynyl, alkoxyalkyl, alkoxyalkoxyalkyl, alkylthioalkyl, alkylthioalkylthioalkyl, alkylthioalkoxyalkyl, alkoxyalkylthioalkyl, carbocyclyl, carbocyclylalkyl, carbocyclylalkoxyalkyl, heterocyclyl, heterocyclylalkyl, and heterocyclylalkoxyalkyl, wherein:

10 any member of such group optionally is substituted with one or more independently selected R^d substituents; and

R^x is alkyl optionally substituted with a substituent selected from the group consisting of R^a-oxy, R^aR^a-amino (wherein each R^a is other than hydrogen), carbocyclyl, and heterocyclyl, wherein:

15 any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, imino, alkoxy, alkylthio, and alkoxyalkoxy, wherein:

20 any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, and alkyl; and

each R^a is independently selected from the group consisting of hydrogen, alkyl, alkoxyalkyl, bisalkoxyalkyl, alkylthioalkyl, alkylsulfoxidoalkyl, alkylsulfonyl, alkylsulfonylalkyl, carbocyclyl, carbocyclylalkyl, carbocyclyloxyalkyl, 25 carbocyclylalkoxyalkyl, carbocyclylthioalkyl, carbocyclylsulfoxidoalkyl, carbocyclylsulfonyl, carbocyclylsulfonylalkyl, heterocyclyl, heterocyclylalkyl, heterocyclyloxyalkyl, heterocyclylalkoxyalkyl, heterocyclylthioalkyl, heterocyclylsulfoxidoalkyl, heterocyclylsulfonyl, and heterocyclylsulfonylalkyl, wherein:

30 any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, imino, alkyl, alkylcarbonyl, carbocyclyl, and carbocyclylalkyl; and
each R^b is independently selected aryl; and

each R^c is independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, -C(H)(NH), -C(H)(NOH), thiol, sulfo, nitro, nitroso, oxo, thioxo, imino, alkyl, alkoxy, alkenyl, alkynyl, alkoxyalkyl, alkylthio, carbocyclyl, carbocyclalkyl, heterocyclyl, and heterocyclalkyl, wherein:

5 any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, imino, alkyl, and carbocyclalkyl; and

10 each R^d is independently selected from the group consisting of halogen, hydroxy, cyano, sulfo, nitro, nitroso, oxo, thioxo, imino, alkyl, alkoxy, alkoxyalkyl, -C(O)(R^e), -S-R^e, -S(O)₂-R^e, carbocyclyl, alkylcarbocyclyl, carbocyclalkyl, heterocyclyl, alkylheterocyclyl, and heterocyclalkyl, wherein:

15 any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino; and

 each R^e is independently selected from the group consisting of hydrogen alkyl, carbocyclyl, carbocyclalkyl, heterocyclyl, and heterocyclalkyl, wherein:

20 any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino; and

 each R^f is independently selected from the group consisting of hydrogen, alkyl, -O-R^h, carbocyclalkyl, and heterocyclalkyl; and



25 any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino; and

 each R^h is independently selected from the group consisting of hydrogen, alkyl, carbocyclyl, carbocyclalkyl, heterocyclyl, and heterocyclalkyl, wherein:



30 any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino.

65. A process according to claim 64, wherein:

the hydroxamic acid compound corresponds in structure to a formula selected from the group consisting of:



5 the unprotected piperidinyll compound corresponds in structure to a formula selected from the group consisting of:




R² is selected from the group consisting of aryl and heteroaryl, wherein:

10 the aryl or heteroaryl optionally is substituted with one or more substituents independently selected from the group consisting of halogen, nitro, nitroso, hydroxy, alkyl, alkoxy, alkylthio, alkoxyalkyl, alkoxyalkylalkyl, C₁-C₂-alkylenedioxy, and alkoxyalkyl, wherein:

15 any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of


halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo,
and imino.

- 5 66. A process according to claim 65, wherein:
the N-alkylating agent is R^x-X^3 ; and
 X^3 is halogen or corresponds in structure to a formula selected from the group
consisting of:



(66-1)

and




(66-2).

- 10 67. A process according to claim 66, wherein X^3 is halogen.
68. A process according to claim 67, wherein contacting of the unprotected
piperidinyll compound with the N-alkylating agent occurs at a temperature of greater than
30°C.
- 15 69. A process according to claim 67, wherein contacting of the unprotected
piperidinyll compound with the N-alkylating agent occurs in the presence of base.
70. A process according to claim 69, wherein the base comprises potassium
carbonate.
- 20 71. A process according to claim 67, wherein:
 X^3 is bromo, and
contacting of the unprotected piperidinyll compound with the N-alkylating agent
occurs at a temperature of from about 50 to about 60°C.
- 25 72. A process according to claim 67, wherein:
 X^3 is chloro, and

contacting of the unprotected piperidiny compound with the N-alkylating agent occurs at a temperature of greater than about 60°C.


73. A process according to claim 67, wherein the hydroxamic acid compound
5 corresponds in structure to Formula (73-1):




74. A process for making an alkylated piperidiny compound or a salt thereof, wherein:

- 10 the process comprises contacting an unprotected piperidiny compound with an N-alkylating agent; and

the alkylated piperidiny compound corresponds in structure to Formula (74-1):



the unprotected piperidiny compound corresponds in structure to Formula (74-2):



15 Z^3 is selected from the group consisting of nitrogen and carbon bonded to hydrogen; and

each n is independently selected from the group consisting of zero, 1, and 2; and

- 20 A is selected from the group consisting of $-\text{O}-$, $-\text{S}-$, $-\text{S}(\text{O})-$, $-\text{S}(\text{O})_2-$, $-\text{C}(\text{O})-$, $-\text{NR}^b-$, $-\text{CO}-\text{N}(\text{R}^b)-$, $-\text{N}(\text{R}^b)-\text{C}(\text{O})-$, $-\text{C}(\text{O})-\text{O}-$, $-\text{O}-\text{C}(\text{O})-$, $-\text{O}-\text{C}(\text{O})-\text{O}-$, $-\text{HC}=\text{CH}-$, $-\text{C}\equiv\text{C}-$, $-\text{N}=\text{N}-$, $-\text{C}(\text{S})-\text{N}(\text{R}^b)-$, $-\text{N}(\text{R}^b)-\text{C}(\text{S})-$, alkyl, alkoxy, oxyalkyl, alkylthio, thioalkyl, and a bond; and

R is selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxyalkyl, carbocyclyl, heterocyclyl, carbocyclylalkyl, heterocyclylalkyl, carbocycloxyalkyl, heterocycloxyalkyl, carbocyclylthioalkyl, and heterocyclylthioalkyl, wherein:

5 any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, nitro, nitroso, hydroxy, oxo, alkyl, alkoxy, alkylthio, alkoxyalkyl, alkoxycarbonylalkyl, C₁-C₂-alkylenedioxy, and alkoxycarbonyl, wherein:

10 any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino; and

E is selected from the group consisting of -O-, -C(O)-, -C(O)-O-, -O-C(O)-, -N(R^b)-, -C(O)-N(R^b)-, -N(R^b)-C(O)-, -C(O)-N(R^b)-N(R^b)-C(O)-, -N(R^b)-C(O)-N(R^b)-, -S-, -S(O)-, -S(O)₂-, -N(R^b)-S(O)₂-, -S(O)₂-N(R^b)-, -O-S(O)₂-, -S(O)₂-O-, -C(NH)-, 15 -C(NOH)-, -N(R^b)-C(NH)-, -N(R^b)-C(NOH)-, -C(NH)-N(R^b)-, -C(NOH)-N(R^b)-, alkyl, alkenyl, carbonylalkyl, alkylcarbonyl, and a bond, wherein:

any alkyl or alkenyl portion of a substituent in such group optionally is substituted with one or more independently selected R^c substituents; and

20 Y is selected from the group consisting of hydrogen, halogen, cyano, alkyl, alkenyl, alkynyl, alkoxyalkyl, alkoxyalkoxyalkyl, alkylthioalkyl, alkylthioalkylthioalkyl, alkylthioalkoxyalkyl, alkoxyalkylthioalkyl, carbocyclyl, carbocyclylalkyl, carbocyclylalkoxyalkyl, heterocyclyl, heterocyclylalkyl, and heterocyclylalkoxyalkyl, wherein:

25 any member of such group optionally is substituted with one or more independently selected R^d substituents; and

R^x is alkyl optionally substituted with a substituent selected from the group consisting of R^a-oxy, R^aR^a-amino (wherein each R^a is other than hydrogen), carbocyclyl, and heterocyclyl, wherein:

30 any member of such group optionally is substituted on any atom capable of such substituent with one or more substituents independently selected from the group consisting of halogen, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, imino, alkoxy, alkylthio, and alkoxyalkoxy, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, and alkyl; and

- each R^a is independently selected from the group consisting of hydrogen, alkyl, alkoxyalkyl, bisalkoxyalkyl, alkylthioalkyl, alkylsulfoxidoalkyl, alkylsulfonyl, alkylsulfonylalkyl, carbocyclyl, carbocyclylalkyl, carbocyclyloxyalkyl, carbocyclylalkoxyalkyl, carbocyclylthioalkyl, carbocyclylsulfoxidoalkyl, carbocyclylsulfonyl, carbocyclylsulfonylalkyl, heterocyclyl, heterocyclylalkyl, heterocyclyloxyalkyl, heterocyclylalkoxyalkyl, heterocyclylthioalkyl, heterocyclylsulfoxidoalkyl, heterocyclylsulfonyl, and heterocyclylsulfonylalkyl, wherein:
- any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, imino, alkyl, alkylcarbonyl, carbocyclyl, and carbocyclylalkyl; and
- each R^b is independently selected aryl; and
- each R^c is independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, -C(H)(NH), -C(H)(NOH), thiol, sulfo, nitro, nitroso, oxo, thioxo, imino, alkyl, alkoxy, alkenyl, alkynyl, alkoxyalkyl, alkylthio, carbocyclyl, carbocyclylalkyl, heterocyclyl, and heterocyclylalkyl, wherein:
- any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, imino, alkyl, and carbocyclylalkyl; and
- each R^d is independently selected from the group consisting of halogen, hydroxy, cyano, sulfo, nitro, nitroso, oxo, thioxo, imino, alkyl, alkoxy, alkoxyalkyl, -C(O)(R^e), -S-R^e, -S(O)₂-R^e, carbocyclyl, alkylcarbocyclyl, carbocyclylalkyl, heterocyclyl, alkylheterocyclyl, and heterocyclylalkyl, wherein:
- any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino; and
- each R^e is independently selected from the group consisting of hydrogen alkyl, carbocyclyl, carbocyclylalkyl, heterocyclyl, and heterocyclylalkyl, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino; and each R^g is independently selected from the group consisting of hydrogen, alkyl, -


5 O- R^h , carbocyclalkyl, and heterocyclalkyl; and

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino; and

each R^h is independently selected from the group consisting of hydrogen, alkyl, carbocycl, carbocyclalkyl, heterocycl, and heterocyclalkyl, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino.


15 75. A process according to claim 74, wherein the alkylated piperidinyl compound corresponds in structure to Formula (75-1):




20 76. A process for making a hydroxamic acid compound or a salt thereof, wherein:

the process comprises contacting a sulfuramine compound with a base to form an anion, and contacting the anion with a carbon dioxide source; and

the hydroxamic acid compound corresponds in structure to Formula (76-1):



the sulfuramine compound corresponds in structure to Formula (76-2):



each n is independently selected from the group consisting of zero, 1, and 2; and
Z is selected from the group consisting of -O-, -S-, -S(O)-, -S(O)₂-, and -N(R^x)-;

5 and

Z³ is selected from the group consisting of nitrogen and carbon bonded to
hydrogen; and

R^x is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, R^a-
oxyalkyl, alkylsulfonyl, aminosulfonyl, R^aR^a-aminoalkyl, carbocyclyl, carbocyclylalkyl,
10 carbocyclylsulfonyl, heterocyclyl, heterocyclylalkyl, and heterocyclylsulfonyl, wherein:
any member of such group optionally is substituted with one or more
substituents independently selected from the group consisting of halogen, hydroxy,
cyano, amino, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, imino, alkyl,
alkoxy, alkylthio, alkoxyalkyl, and alkoxyalkoxy, wherein:

15 any member of such group optionally is substituted with one or
more substituents independently selected from the group consisting of
halogen, hydroxy, and alkyl; and

A is selected from the group consisting of -O-, -S-, -S(O)-, -S(O)₂-, -C(O)-, -NR^b-,
-CO-N(R^b), -N(R^b)-C(O)-, -C(O)-O-, -O-C(O)-, -O-C(O)-O-, -HC=CH-, -C≡C-, -N=N-,
20 -C(S)-N(R^b)-, -N(R^b)-C(S)-, alkyl, alkoxy, oxyalkyl, alkylthio, thioalkyl, and a bond; and

R is selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxyalkyl,
carbocyclyl, heterocyclyl, carbocyclylalkyl, heterocyclylalkyl, carbocycliloxyalkyl,
heterocycliloxyalkyl, carbocyclylthioalkyl, and heterocyclylthioalkyl, wherein:

25 any member of such group optionally is substituted with one or more
substituents independently selected from the group consisting of halogen, nitro,
nitroso, hydroxy, oxo, alkyl, alkoxy, alkylthio, alkoxyalkyl, alkoxyalkylalkyl,
C₁-C₂-alkylenedioxy, and alkoxyalkyl, wherein:

any member of such group optionally is substituted with one or
more substituents independently selected from the group consisting of

halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino; and

E is selected from the group consisting of -O-, -C(O)-, -C(O)-O-, -O-C(O)-, -N(R^b)-, -C(O)-N(R^b)-, -N(R^b)-C(O)-, -C(O)-N(R^b)-N(R^b)-C(O)-, -N(R^b)-C(O)-N(R^b)-, -S-, -S(O)-, -S(O)₂-, -N(R^b)-S(O)₂-, -S(O)₂-N(R^b)-, -O-S(O)₂-, -S(O)₂-O-, -C(NH)-, -C(NOH)-, -N(R^b)-C(NH)-, -N(R^b)-C(NOH)-, -C(NH)-N(R^b)-, -C(NOH)-N(R^b)-, alkyl, alkenyl, carbonylalkyl, alkylcarbonyl, and a bond, wherein:

any alkyl or alkenyl portion of a substituent in such group optionally is substituted with one or more independently selected R^c substituents; and

Y is selected from the group consisting of hydrogen, halogen, cyano, alkyl, alkenyl, alkynyl, alkoxyalkyl, alkoxyalkoxyalkyl, alkylthioalkyl, alkylthioalkylthioalkyl, alkylthioalkoxyalkyl, alkoxyalkylthioalkyl, carbocyclyl, carbocyclylalkyl, carbocyclylalkoxyalkyl, heterocyclyl, heterocyclylalkyl, and heterocyclylalkoxyalkyl, wherein:

any member of such group optionally is substituted with one or more independently selected R^d substituents; and

each R^a is independently selected from the group consisting of hydrogen, hydroxy, alkyl, alkenyl, alkynyl, alkoxy, alkoxyalkyl, bisalkoxyalkyl, alkylthioalkyl, alkylthioalkenyl, alkylsulfoxidoalkyl, alkylsulfonyl, alkylsulfonylalkyl, carbocyclyl, carbocyclylalkyl, carbocyclylalkoxyalkyl, carbocyclylthioalkyl, carbocyclylthioalkenyl, carbocyclylsulfoxidoalkyl, carbocyclylsulfonyl, carbocyclylsulfonylalkyl, heterocyclyl, heterocyclylalkyl, heterocyclylalkoxyalkyl, heterocyclylthioalkyl, heterocyclylsulfoxidoalkyl, heterocyclylsulfonyl, heterocyclylsulfonylalkyl, aminoalkyl, aminosulfonyl, aminoalkylsulfonyl, and alkoxyalkylaminoalkyl, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, imino, alkyl, alkylcarbonyl, carbocyclyl, and carbocyclylalkyl; and

each R^b is independently selected aryl; and

each R^c is independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, -C(H)(NH), -C(H)(NOH), thiol, sulfo, nitro, nitroso, oxo, thioxo, imino,

alkyl, alkoxy, alkenyl, alkynyl, alkoxyalkyl, alkylthio, carbocyclyl, carbocyclylalkyl, heterocyclyl, and heterocyclylalkyl, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, imino, alkyl, and carbocyclylalkyl; and

each R^d is independently selected from the group consisting of halogen, hydroxy, cyano, sulfo, nitro, nitroso, oxo, thioxo, imino, alkyl, alkoxy, alkoxyalkyl, -C(O)(R^g), -S-R^e, -S(O)₂-R^e, carbocyclyl, alkylcarbocyclyl, carbocyclylalkyl, heterocyclyl,

alkylheterocyclyl, and heterocyclylalkyl, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino; and

each R^e is independently selected from the group consisting of hydrogen alkyl, carbocyclyl, carbocyclylalkyl, heterocyclyl, and heterocyclylalkyl, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino; and

each R^g is independently selected from the group consisting of hydrogen, alkyl, -O-R^h, carbocyclylalkyl, and heterocyclylalkyl; and



any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino; and

each R^h is independently selected from the group consisting of hydrogen, alkyl, carbocyclyl, carbocyclylalkyl, heterocyclyl, and heterocyclylalkyl, wherein:



any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino.

77. A process according to claim 76, wherein:

the hydroxamic acid compound corresponds in structure to a formula selected from the group consisting of:



the sulfuramine compound corresponds in structure to a formula selected from the group consisting of:



R^2 is selected from the group consisting of aryl and heteroaryl, wherein:

the aryl or heteroaryl optionally is substituted with one or more substituents independently selected from the group consisting of halogen, nitro, nitroso, hydroxy, alkyl, alkoxy, alkylthio, alkoxyalkyl, alkoxyalkylalkyl, C_1 - C_2 -alkylenedioxy, and alkoxyalkyl, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino.

78. A process according to claim 77, wherein the base comprises a strong, non-aqueous base.

79. A process according to claim 78, wherein the base comprises alkyllithium.

80. A process according to claim 79, wherein the base comprises butyllithium.

81. A process according to claim 78, wherein the base comprises lithium diisopropylamide.

5


82. A process according to claim 78, wherein contacting of the sulfuramine compound with the base occurs at a temperature of less than 20°C.

83. A process according to claim 78, wherein contacting of the anion with the
10 carbon dioxide source occurs at a temperature of less than 20°C.

84. A process according to claim 78, wherein the source of carbon dioxide comprises methylchloroformate.

15 85. A process according to claim 78, wherein the source of carbon dioxide comprises CO₂ gas.

86. A process according to claim 85, wherein the sulfuramine compound corresponds in structure to Formula (86-1):




20

(86-1).

87. A process for making a carboxylic acid compound, wherein:


the process comprises contacting a sulfuramine compound with a base to form an anion, and contacting the anion with a carbon dioxide source; and

25 the carboxylic acid compound corresponds in structure to Formula (87-1):



(87-1); and

the sulfuramine compound corresponds in structure to Formula (87-2):



(87-2); and

each n is independently selected from the group consisting of zero, 1, and 2; and

Z is selected from the group consisting of -O-, -S-, -S(O)-, -S(O)₂-, and -N(R^x)-;

and

Z³ is selected from the group consisting of nitrogen and carbon bonded to hydrogen; and

R^x is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, R^a-oxyalkyl, alkylsulfonyl, aminosulfonyl, R^aR^a-aminoalkyl, carbocyclyl, carbocyclylalkyl, carbocyclylsulfonyl, heterocyclyl, heterocyclylalkyl, and heterocyclylsulfonyl, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, amino, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, imino, alkyl, alkoxy, alkylthio, alkoxyalkyl, and alkoxyalkoxy, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, and alkyl; and

A is selected from the group consisting of -O-, -S-, -S(O)-, -S(O)₂-, -C(O)-, -NR^b-, -CO-N(R^b), -N(R^b)-C(O)-, -C(O)-O-, -O-C(O)-, -O-C(O)-O-, -HC=CH-, -C≡C-, -N=N-, -C(S)-N(R^b)-, -N(R^b)-C(S)-, alkyl, alkoxy, oxyalkyl, alkylthio, thioalkyl, and a bond; and

R is selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxyalkyl, carbocyclyl, heterocyclyl, carbocyclylalkyl, heterocyclylalkyl, carbocycliloxyalkyl, heterocycliloxyalkyl, carbocyclylthioalkyl, and heterocyclylthioalkyl, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, nitro,

nitroso, hydroxy, oxo, alkyl, alkoxy, alkylthio, alkoxyalkyl, alkoxyacetylalkyl,
C₁-C₂-alkylenedioxy, and alkoxyacetyl, wherein:

any member of such group optionally is substituted with one or
more substituents independently selected from the group consisting of
5 halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thiooxo,
and imino; and

E is selected from the group consisting of -O-, -C(O)-, -C(O)-O-, -O-C(O)-,
-N(R^b)-, -C(O)-N(R^b)-, -N(R^b)-C(O)-, -C(O)-N(R^b)-N(R^b)-C(O)-, -N(R^b)-C(O)-N(R^b)-,
-S-, -S(O)-, -S(O)₂-, -N(R^b)-S(O)₂-, -S(O)₂-N(R^b)-, -O-S(O)₂-, -S(O)₂-O-, -C(NH)-,
10 -C(NOH)-, -N(R^b)-C(NH)-, -N(R^b)-C(NOH)-, -C(NH)-N(R^b)-, -C(NOH)-N(R^b)-, alkyl,
alkenyl, carbonylalkyl, alkylcarbonyl, and a bond, wherein:

any alkyl or alkenyl portion of a substituent in such group optionally is
substituted with one or more independently selected R^c substituents; and

Y is selected from the group consisting of hydrogen, halogen, cyano, alkyl,
15 alkenyl, alkynyl, alkoxyalkyl, alkoxyalkoxyalkyl, alkylthioalkyl, alkylthioalkylthioalkyl,
alkylthioalkoxyalkyl, alkoxyalkylthioalkyl, carbocyclyl, carbocyclylalkyl,
carbocyclylalkoxyalkyl, heterocyclyl, heterocyclylalkyl, and heterocyclylalkoxyalkyl,
wherein:

any member of such group optionally is substituted with one or more
20 independently selected R^d substituents; and

each R^a is independently selected from the group consisting of hydrogen, hydroxy,
alkyl, alkenyl, alkynyl, alkoxy, alkoxyalkyl, bisalkoxyalkyl, alkylthioalkyl,
alkylthioalkenyl, alkylsulfoxidoalkyl, alkylsulfonyl, alkylsulfonylalkyl, carbocyclyl,
carbocyclylalkyl, carbocyclylalkoxyalkyl, carbocyclylthioalkyl,
25 carbocyclylthioalkenyl, carbocyclylsulfoxidoalkyl, carbocyclylsulfonyl,
carbocyclylsulfonylalkyl, heterocyclyl, heterocyclylalkyl, heterocyclylalkoxyalkyl,
heterocyclylthioalkyl, heterocyclylsulfoxidoalkyl,
heterocyclylsulfonyl, heterocyclylsulfonylalkyl, aminoalkyl, aminosulfonyl,
aminoalkylsulfonyl, and alkoxyalkylaminoalkyl, wherein:

30 any member of such group optionally is substituted with one or more
substituents independently selected from the group consisting of halogen, hydroxy,
cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thiooxo, imino, alkyl, alkylcarbonyl,
carbocyclyl, and carbocyclylalkyl; and

each R^b is independently selected aryl; and

each R^c is independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, -C(H)(NH), -C(H)(NOH), thiol, sulfo, nitro, nitroso, oxo, thioxo, imino, alkyl, alkoxy, alkenyl, alkynyl, alkoxyalkyl, alkylthio, carbocyclyl, carbocyclylalkyl,

5 heterocyclyl, and heterocyclylalkyl, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, imino, alkyl, and carbocyclylalkyl; and

10 each R^d is independently selected from the group consisting of halogen, hydroxy, cyano, sulfo, nitro, nitroso, oxo, thioxo, imino, alkyl, alkoxy, alkoxyalkyl, -C(O)(R^g), -S-R^e, -S(O)₂-R^e, carbocyclyl, alkylcarbocyclyl, carbocyclylalkyl, heterocyclyl, alkylheterocyclyl, and heterocyclylalkyl, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino; and

15 each R^e is independently selected from the group consisting of hydrogen alkyl, carbocyclyl, carbocyclylalkyl, heterocyclyl, and heterocyclylalkyl, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino; and

20 each R^g is independently selected from the group consisting of hydrogen, alkyl, -O-R^h, carbocyclylalkyl, and heterocyclylalkyl; and


any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino; and

25 each R^h is independently selected from the group consisting of hydrogen, alkyl, carbocyclyl, carbocyclylalkyl, heterocyclyl, and heterocyclylalkyl, wherein:


any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino.

30

88. A process according to claim 87, wherein the carboxylic acid compound corresponds in structure to Formula (88-1):



89. A compound or a salt thereof, wherein:
the compound corresponds in structure to Formula (89-1):



n is selected from the group consisting of zero, 1, and 2; and

Z^1 is selected from the group consisting of $-N(H)-$, X^1 , and $-N(R^{x2})-$; and

Z^3 is selected from the group consisting of nitrogen and carbon bonded to hydrogen; and

X^1 is selected from the group consisting of $-O-$, $-S-$, $-S(O)-$, $-S(O)_2-$, and $-N(R^{x1})-$; and

R^{x1} is a nitrogen-protecting group; and

R^{x2} is alkyl optionally substituted with a substituent selected from the group consisting of R^a -oxy, R^aR^a -amino (wherein each R^a is other than hydrogen), carbocyclyl, and heterocyclyl, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, imino, alkoxy, alkylthio, and alkoxyalkoxy, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, and alkyl; and

A is selected from the group consisting of -O-, -S-, -S(O)-, -S(O)₂-, -C(O)-, -NR^b-, -CO-N(R^b), -N(R^b)-C(O)-, -C(O)-O-, -O-C(O)-, -O-C(O)-O-, -HC=CH-, -C≡C-, -N=N-, -C(S)-N(R^b)-, -N(R^b)-C(S)-, alkyl, alkoxy, oxyalkyl, alkylthio, thioalkyl, and a bond; and

R is selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxyalkyl, carbocyclyl, heterocyclyl, carbocyclylalkyl, heterocyclylalkyl, carbocycloxyalkyl, heterocycloxyalkyl, carbocyclylthioalkyl, and heterocyclylthioalkyl, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, nitro, nitroso, hydroxy, oxo, alkyl, alkoxy, alkylthio, alkoxyalkyl, alkoxycarbonylalkyl, C₁-C₂-alkylenedioxy, and alkoxycarbonyl, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino; and

E is selected from the group consisting of -O-, -C(O)-, -C(O)-O-, -O-C(O)-, -N(R^b)-, -C(O)-N(R^b)-, -N(R^b)-C(O)-, -C(O)-N(R^b)-N(R^b)-C(O)-, -N(R^b)-C(O)-N(R^b)-, -S-, -S(O)-, -S(O)₂-, -N(R^b)-S(O)₂-, -S(O)₂-N(R^b)-, -O-S(O)₂-, -S(O)₂-O-, -C(NH)-, -C(NOH)-, -N(R^b)-C(NH)-, -N(R^b)-C(NOH)-, -C(NH)-N(R^b)-, -C(NOH)-N(R^b)-, alkyl, alkenyl, carbonylalkyl, alkylcarbonyl, and a bond, wherein:

any alkyl or alkenyl portion of a substituent in such group optionally is substituted with one or more independently selected R^c substituents; and

Y is selected from the group consisting of hydrogen, halogen, cyano, alkyl, alkenyl, alkynyl, alkoxyalkyl, alkoxyalkoxyalkyl, alkylthioalkyl, alkylthioalkylthioalkyl, alkylthioalkoxyalkyl, alkoxyalkylthioalkyl, carbocyclyl, carbocyclylalkyl, carbocyclylalkoxyalkyl, heterocyclyl, heterocyclylalkyl, and heterocyclylalkoxyalkyl, wherein:

any member of such group optionally is substituted with one or more independently selected R^d substituents; and

each R^a is independently selected from the group consisting of hydrogen, alkyl, alkoxyalkyl, bisalkoxyalkyl, alkylthioalkyl, alkylsulfoxidoalkyl, alkylsulfonyl, alkylsulfonylalkyl, carbocyclyl, carbocyclylalkyl, carbocycloxyalkyl, carbocyclylalkoxyalkyl, carbocyclylthioalkyl, carbocyclylsulfoxidoalkyl,

carbocyclylsulfonyl, carbocyclylsulfonylalkyl, heterocyclyl, heterocyclylalkyl, heterocyclyloxyalkyl, heterocyclylalkoxyalkyl, heterocyclylthioalkyl, heterocyclylsulfoxidoalkyl, heterocyclylsulfonyl, and heterocyclylsulfonylalkyl, wherein:

any member of such group optionally is substituted with one or more

substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, imino, alkyl, alkylcarbonyl, carbocyclyl, and carbocyclylalkyl; and

each R^b is independently selected aryl; and

each R^c is independently selected from the group consisting of halogen, hydroxy,

cyano, carboxy, -C(H)(NH), -C(H)(NOH), thiol, sulfo, nitro, nitroso, oxo, thioxo, imino, alkyl, alkoxy, alkenyl, alkynyl, alkoxyalkyl, alkylthio, carbocyclyl, carbocyclylalkyl, heterocyclyl, and heterocyclylalkyl, wherein:

any member of such group optionally is substituted with one or more

substituents independently selected from the group consisting of halogen, hydroxy,

cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, imino, alkyl, and

carbocyclylalkyl; and

each R^d is independently selected from the group consisting of halogen, hydroxy,

cyano, sulfo, nitro, nitroso, oxo, thioxo, imino, alkyl, alkoxy, alkoxyalkyl, -C(O)(R^g), -S-R^e, -S(O)₂-R^e, carbocyclyl, alkylcarbocyclyl, carbocyclylalkyl, heterocyclyl,

alkylheterocyclyl, and heterocyclylalkyl, wherein:

any member of such group optionally is substituted with one or more

substituents independently selected from the group consisting of halogen, hydroxy,

cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino; and

each R^e is independently selected from the group consisting of hydrogen alkyl,

carbocyclyl, carbocyclylalkyl, heterocyclyl, and heterocyclylalkyl, wherein:

any member of such group optionally is substituted with one or more

substituents independently selected from the group consisting of halogen, hydroxy,

cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino; and

each R^g is independently selected from the group consisting of hydrogen, alkyl, -

O-R^h, carbocyclylalkyl, and heterocyclylalkyl; and

any member of such group optionally is substituted with one or more

substituents independently selected from the group consisting of halogen, hydroxy,


cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino; and

each R^h is independently selected from the group consisting of hydrogen, alkyl, carbocyclyl, carbocyclylalkyl, heterocyclyl, and heterocyclylalkyl, wherein:

any member of such group optionally is substituted with one or more substituents independently selected from the group consisting of halogen, hydroxy, cyano, carboxy, thiol, sulfo, nitro, nitroso, oxo, thioxo, and imino.

90. A compound or salt thereof according to claim 89, wherein Z¹ is -N(H)-.

91. A compound or salt thereof according to claim 90, wherein the compound corresponds in structure to Formula (91-1):





92. A compound or salt thereof according to claim 89, wherein Z¹ is X¹.


93. A compound or salt thereof according to claim 92, wherein:

X¹ is -N(R^{x1})-; and

R^{x1} is selected from the group consisting of alkoxyalkyl, alkoxycarbonyl, and arylalkoxycarbonyl.

94. A compound or salt thereof according to claim 93, wherein the compound corresponds is selected from the group consisting of:





(94-3).